

# Effect of industrialization on allergic diseases in school children

Ahmet Ozdemir<sup>1</sup>, Demet Can<sup>1</sup>, Ilker Gunay<sup>1</sup>, Tekin Nacaroglu<sup>1</sup>, Canan Sule Karkiner<sup>1</sup>, Ala Ustyo<sup>1</sup>, Hacer Kamali<sup>1</sup>, Muge Ayanoglu<sup>1</sup>, Turkan Gunay<sup>2</sup>, Done Dogan<sup>1</sup>

<sup>1</sup>Dr. Behcet Uz Children's Hospital Department of Pediatric Allergy, Izmir, Turkey

<sup>2</sup>Dokuz Eylul University Faculty of Medicine, Department of Public Health, Izmir, Turkey

Copyright © 2018 by authors and Annals of Medical Research Publishing Inc.

## Abstract

**Aim:** The lifestyle and environmental changes associated with industrialization and urbanization are crucial factors in determining the prevalence of allergic diseases.

**Material and Methods:** The aim of this study was to determine the prevalence of asthma and other allergic disorders in an industrial town. Study Design: Prospective, Cross-sectional study. This cross-sectional population-based study was conducted in Kemalpaşa, Izmir, Turkey, which is an industrial town. Using the International Study of Asthma and Allergies in Childhood (ISAAC), the questionnaire was carried out on 1511 school-children aged between 13-14 years living in the district center.

**Results:** The ISAAC Phase One questionnaire was answered by 1373 (90.87%) students. Six hundred ninety two (50.5%) of the children were girls and 681 (49.5%) were boys. The frequency of ever wheezing was 34%, that of current wheezing was 31.1% and that of physician-diagnosed asthma was 4.1%. The frequency of ever rhinitis was 36.6%, current rhinitis was 31.3%, current allergic rhino-conjunctivitis was 22.1% and physician-diagnosed allergic rhinitis was 11.1%. The frequency of ever eczema was 11.3% and that of physician-diagnosed eczema was 3.6%.

**Conclusion:** The prevalence of asthma was found to be significantly higher, although there was no significant difference in allergic rhinitis and eczema compared with non-industrial neighboring states. The incidence of asthma was higher than other provinces which may be explained by the risk factors associated with this industrialized area. The fact that only children of the same age group and residence were evaluated contributed to the epidemiological value of our study.

**Keywords:** Allergic Rhinitis; Asthma; Atopic Dermatitis; Children; ISAAC; Industrial Region.

## INTRODUCTION

The International Study of Asthma and Allergies in Childhood Phase Three (ISAAC-III) is used to examine the changes in the prevalence of symptoms of allergic disorders. Most of the ISAAC-III studies revealed an increase in the burden of allergic diseases in developing countries (1).

In ISAAC-III, risk factors had been explored using an environmental questionnaire. Many factors, including paracetamol, antibiotics, and truck traffic exposure have been explored (2-4). Environmental changes, western style, living conditions, and rapid industrialization without proper planning may play a role in the increased prevalence of allergic symptoms (5-7) although this has not been proven yet.

In this study, we aimed to determine the prevalence of asthma and other allergic diseases in an industrial area.

In order to avoid randomization bias, only children in a certain area were included in the study group.

## MATERIAL and METHODS

### Study Population

Our study was done in the Kemalpaşa district of Izmir, Turkey. Kemalpaşa is the major industrial region of Izmir, which is the third largest city in Turkey. There are 5 elementary schools in the district center. The total number of students aged 13-14 was 1511. All students in the target age group were included in the study.

### Questionnaires and Definitions

First, researchers gave a brief presentation on allergic diseases to all teachers and students. Then all students were given the ISAAC Phase One questionnaire. The questionnaires were completed by the children themselves. The severity of asthma symptoms was assessed by questions dealing with the last 12 months. Symptoms

Received: 14.03.2018 Accepted: 20.03.2018 Available online:

Corresponding Author: Ahmet Ozdemir, Dr. Behcet Uz Children's Hospital Department of Pediatric Allergy, Izmir, Turkey  
E-mail: dr.ozdemir@yahoo.com

that were asked about included number of wheezing attacks, sleep disturbance due to wheezing, and severe wheezing that limits speech to only one or two words at a time between breaths. If a student answered the question "Have you had wheezing or whistling in the chest in the past 12 months?" as "yes", then s/he was considered to have asthma (current wheezing). If a student answered the question "Have you ever had wheezing or whistling in the chest?" as "yes", then s/he was considered to have asthma (ever wheezing). "Ever rhinitis" was defined as a positive response to, "Have you ever had a problem with sneezing or a runny or blocked nose when you (he/she) did not have a cold or the flu?". "Current rhinitis" was defined as a positive response to "In the past 12 months, has your child had a problem with sneezing or a runny or blocked nose when he/she did not have a cold or the flu?". Patients with symptoms occurring only between March and October were classified as having "seasonal rhinitis". When the symptoms were reported as occurring beyond the March–October period, the classification was "perennial rhinitis." Current allergic rhino conjunctivitis symptoms were evaluated as "positive" if both of the following questions were answered by the parent as "yes": "In the past 12 months, has your child had a problem with sneezing or a runny or blocked nose when he/she did not have a cold or the flu?" and "In the past 12 months, has this nose problem been accompanied by itchy watery eyes?" Similarly, symptoms for current eczema were studied on the basis of positive answers to two questions: "Has your child had this itchy rash at any time in the past 12 months?" and "Has this itchy rash at any time affected any of the following places: folds of the elbows, behind the knees, in front of the ankles, under the buttocks, or around the neck, ears, or eyes."

#### Ethical Considerations

Permission for the study was obtained from the local ethics committee, and the central and provincial directors of the Ministry of Education and town governors. Written consent was obtained from each participant.

#### Statistical Analysis

The SPSS-15 statistical software package (SPSS, Inc., Chicago, IL) was used for the analysis. Statistical analysis included frequency and percent distributions, calculation of prevalence rates for allergic diseases, and comparisons using chi-square and Student's t-tests.

## RESULTS

1373 (90.8%) students were given the ISAAC Phase One Questionnaire. None of them declined to participate in the study. Questionnaires could not be given to those students who were not at school on the day of data collection. Of the 1373 students, 47.8% had an allergic symptom. The prevalence of asthma, rhinitis and eczema was 34%, 37% and 11%, respectively.

#### Gender

There was a slight majority of girls (698, 50.5%). Allergic symptoms were seen in girls more frequently (58.4%) than boys. This difference was significant ( $p < 0.05$ ). In the asthma group, 60% of the children were girls. Girls represented 65% of the rhinitis and 69% of the eczema groups. Gender difference for all groups was significant ( $p < 0.05$ ). In case of co-existing allergic problems, the percentage of girls was significantly more than that of

boys ( $p < 0.05$ ).

#### Asthma

The frequency of symptoms related to asthma is shown in Table 1. The ISAAC Phase One questionnaire revealed ever wheezing (EW) in 34%, and current wheezing (CW) in 31.1% of the study group. In both wheezing groups, there were significantly more girls than boys ( $p < 0.05$ ).

**Table 1. Frequency of asthma-related findings**

Findings	n (%)
Ever Wheezing	468 (34)
Current Wheezing	428 (31.1)
Number of attacks	
1-3	67 (4.8)
4-12	24 (1.7)
> 12	68 (4.9)
Wheeze disturbs sleep	
Never	89 (6.4)
Less than once a week	42 (3.0)
More than once a week	29 (2.1)
Severe wheeze limiting speech	117 (8.5)
Physician-diagnosed asthma	57 (4.1)
Physician-diagnosed allergic bronchitis	168 (12.2)
Exercise wheeze	256 (18.6)
Night cough	278 (20.2)

#### Rhinitis

The frequency of ever rhinitis was 36.6%, current rhinitis was 31.3%, current allergic rhino-conjunctivitis was 22.1%, physician-diagnosed allergic rhinitis was 11.1%, seasonal rhinitis was 8.8% and perennial rhinitis was 1.3%. Both in the seasonal and perennial rhinitis groups, girls represented more than 60% of the group.

#### Eczema

The frequency of ever eczema was 11.3% and that of physician-diagnosed eczema was 3.6% (Table 2).

**Table 2. Frequency of eczema-related findings**

Finding	n (%)
Ever eczema	156 (11.3)
Current eczema	81 (5.8)
Site of itchy rash	
Folds of the elbows	37 (2.6)
Behind the knees	33 (2.4)
In front of the ankles	30 (2.2)
Neck	40 (2.9)
Around the ears	15 (1.1)
Around the eyes	21 (1.5)
Rash disappearing during the last 12 months	74 (5.4)
Sleeplessness due to rash during the last 12 months	
Never	64 (4.6)
< one night a week	25 (1.8)
≥ one night a week	13 (0.9)
Physician-diagnosed eczema	50 (3.6)

## DISCUSSION

The prevalence of rhinitis and eczema in our study was consistent with other ISAAC studies that were done in the same region (8). However, asthma prevalence was found to be higher. This may be due to the rapid urbanization

and industrialization of the Kemalpaşa area.

Before SAAC different centers had used different methods so the findings were not comparable between centers. To counter this problem, ISAAC formulated a questionnaire for the purpose of allowing international comparisons of asthma prevalence and severity. For this reason ISAAC is very important and is widely utilized (7-10).

In Turkey, the first ISAAC Phase One study was done in 1995 and CW prevalence was found to be 9.8% and EW 15.1% (11). In 1999, the first ISAAC Phase Two study revealed 11.5% CW and 6.9% physician-diagnosed asthma (12). During the same period, EW prevalence was found to be 21.2%, asthma 6.4%, rhinitis 13.6% and eczema 23.7% in the Aegean region, in which Kemalpaşa is located (8). A number of studies have found similar figures in this region. CW prevalence was between 14.7-15.9%, rhinitis prevalence was between 14.5-46% and that of eczema was between 7.2-9-6% (13-15). Although EW, rhinitis and eczema were as frequent as in the other studies in the same region, CW was far more prevalent, as high as two fold.

The question here is what was the reason behind the high prevalence of CW?" It was probably not of genetic origin, because Kemalpaşa has a very cosmopolitan population, due to internal migration. This is what makes us think that environmental factors are the cause of this high prevalence of CW. Being an area of heavy industry results in migration, crowded households air pollution and low income.

While allergic diseases have had a tendency to increase in industrialized countries, the rate of increase has been slowing down recently (16-18).

The first ISAAC Phase Three study in Turkey revealed an increase in the prevalence of both EW (from 15.1% to 25.3%) and CW (from 8.2% to 11.3%). Family history of asthma, eczema and food allergy diagnosed by a physician, were associated significantly with high risk for asthma (19). Unlike this study, our study revealed the importance of environmental factors, as in some other developing countries (20).

Since Kemalpaşa is an industrial area, there has been rapid urbanization in the recent years. The lifestyle and environmental changes associated with urbanization are crucial factors in determining the prevalence of allergic diseases in developing countries. For example, in China, marked differences have been observed in the prevalence of asthma between rural and urban communities. The prevalence of asthma symptoms in adolescents living in urban Beijing was six- to ten fold greater than that of outlying rural communities (21). Similar findings have been observed in another study conducted in Africa, in which the rate of exercise-induced asthma was 50-fold higher in urban communities than in rural ones (22).

When we evaluate asthma data, both the number and the severity of the attacks were high (Table 1). In developed countries, although there is an increasing tendency for current wheezing prevalence, the prevalence of severity is decreasing. On the other hand, there is an increase in the

severity of symptoms in developing countries (23).

There is a growing amount of evidence that air pollution may aggravate asthma. The level of particles, however, could not explain the regional differences (24). The high frequency of allergic diseases in our study may be due to air pollution, since Kemalpaşa is a dense industrial area. There are studies showing the relationship between air pollution and asthma prevalence (25,26).

Although air pollution has not been shown to be a risk factor by ISAAC studies, truck traffic exposure was described as a risk factor in developed countries. ISAAC Phase Three revealed a positive association among truck traffic density on the street of residency and asthma, rhino-conjunctivitis and eczema symptom frequencies (4). Kemalpaşa also has traffic pollution, but we did not have any regular air pollution data. This was one of the major limitations of our study.

All allergic diseases were more frequent in girls. Similar findings were reported from Israel (7). Asthma symptoms were reported to be more frequent among females in a study involving 4 countries: Australia, England, Germany and New Zealand (27). These findings are in accordance with studies showing a male predominance for wheezing during the first decade of life, which is reversed around the time of puberty (28). The high prevalence of allergic diseases in girls may be caused by environmental factors, rather than atopy, in which there is a male dominance.

There are a number of ISAAC studies evaluating the influence of environmental and socioeconomic factors for asthma, allergic rhinitis, and eczema among children living in rural and urban areas. A number of studies have analyzed the differences in asthma prevalence between rural and urban areas with conflicting results (29). In a study which was done in Kenya, environmental factors were found to be associated with the rural-urban difference (30). Allergic symptoms were found more frequently in urban areas in Belarus (31).

## CONCLUSION

In conclusion, we have included the whole population of children in the 13-14 age group in a certain area and were able to determine the prevalence of allergic diseases. When compared with neighboring areas, there were no differences in rhinitis and eczema prevalence; asthma was seen more frequently, and this was considered to be related with the industrialization of the district. Rapid urbanization without an adequate infrastructure is a common problem in Turkey, and Kemalpaşa is a striking example of this situation. There are some environmental factors, the effects of which have not been clearly identified, which require further investigation.

### Acknowledgements

*Competing interests: The authors declare that they have no competing interest.*

*Funding: No financial disclosure was declared by the authors.*

*Ethical approval: This study was Institutional Review Board approved.*

## REFERENCES

1. Asher MI, Montefort S, Björkstén B, Lai CK, Strachan DP, Weiland SK, et al. ISAAC Phase Three Study Group. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* 2006;368(9537):733-43.
2. Beasley R, Clayton T, Crane J, von Mutius E, Lai CK, Montefort S, et al. ISAAC Phase Three Study Group. Association between paracetamol use in infancy and childhood, and risk of asthma, rhinoconjunctivitis, and eczema in children aged 6-7 years: analysis from Phase Three of the ISAAC programme. *Lancet* 2008;372(9643):1039-48.
3. Foliaki S, Pearce N, Björkstén B, Mallol J, Montefort S, von Mutius E. International Study of Asthma and Allergies in Childhood Phase III Study Group. Antibiotic use in infancy and symptoms of asthma, rhinoconjunctivitis, and eczema in children 6 and 7 years old: International Study of Asthma and Allergies in Childhood Phase III. *J Allergy Clin Immunol* 2009;124(5):982-89.
4. Brunekreef B, Stewart AW, Anderson HR, Lai CK, Strachan DP, Pearce N. ISAAC Phase 3 Study Group. Self-reported truck traffic on the street of residence and symptoms of asthma and allergic disease: a global relationship in ISAAC phase 3. *Environ Health Perspect* 2009;117(11):1791-8.
5. Sestini P, De Sario M, Bugiani M, Bisanti L, Giannella G, Kaisermann D, et al. Frequency of asthma and allergies in Italian children and adolescents: results from SIDRIA-2. *Epidemiol Prev* 2005;29(2 Suppl):24-31.
6. Ige OM, Falade AG, Arinola OG. Atopy is a risk factor for adult asthma in urban community of Southwestern Nigeria. *Lung India* 2012;29(2):114-9.
7. Shohat T, Golan G, Tamir R, Green MS, Livne I, Davidson Y, et al. Prevalence of asthma in 13-14 yr-old schoolchildren across Israel. *Eur Respir J* 2000;15(4):725-9.
8. Asher MI, Keil U, Anderson HR, Beasley R, Crane J, Martinez F, et al. International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *Eur Respir J* 1995;8(3):483-91.
9. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. *Lancet* 1998;351(9111):1225-32.
10. Oneş U, Sapan N, Somer A, Dişçi R, Salman N, Güler N, et al. Prevalence of childhood asthma in Istanbul, Turkey. *Allergy* 1997;52(5):570-75.
11. Saraçlar Y, Kuyucu S, Tuncer A, Sekerel B, Saçkesen C, Kocabaş C. Prevalence of asthmatic phenotypes and bronchial hyper responsiveness in Turkish school children: an International Study of Asthma and Allergies in Childhood (ISAAC) phase 2 study. *Ann Allergy Asthma Immunol* 2003;91(5):477-84.
12. Demir E, Tanaç R, Can D, Gülen F, Yenigün A, Aksakal K. Is there an increase in the prevalence of allergic diseases among schoolchildren from the Aegean region of Turkey? *Allergy Asthma Proc* 2005;26(5):410-4.
13. Karaman O, Turgut CS, Uzuner N, Olmez D, Babayigit A, Kose S, et al. The determination of asthma, rhinitis, eczema, and atopy prevalence in 9- to 11-year-old children in the city of Izmir. *Allergy Asthma Proc* 2006;27(4):319-24.
14. Yuksel H, Dinc G, Sakar A, Yilmaz O, Yorgancıoglu A, Celik P, et al. Prevalence and comorbidity of allergic eczema, rhinitis, and asthma in a city in western Turkey. *J Investig Allergol Clin Immunol* 2008;18(1):31-35.
15. Civelek E, Cakir B, Boz AB, Yuksel H, Orhan F, Uner A, et al. Extent and Burden of Allergic Diseases in Elementary Schoolchildren: A National Multicenter Study. *J Investig Allergol Clin Immunol* 2010;20(4):280-8.
16. Ronchetti R, Villa MP, Barreto M, Rota R, Pagani J, Martella S, et al. Is the increase in childhood asthma coming to an end? Findings from three surveys of schoolchildren in Rome, Italy. *Eur Respir J* 2001;17(5):881-6.
17. Grize L, Gassner M, Wuthrich B, Bringolf-Isler B, Takken-Sahli K, Sennhauser FH, et al. Trends in prevalence of asthma, allergic rhinitis and atopic dermatitis in 5-7-year old Swiss children from 1992 to 2001. *Allergy* 2006;61(5):556-62.
18. Pearce N, Ait-Khaled N, Beasley R, Mallol J, Keil U, Mitchell E, et al. Worldwide trends in the prevalence of asthma symptoms: phase III of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax* 2007;62(9):758-66.
19. Ones U, Akcay A, Tamay Z, Guler N, Zencir M. Rising trend of asthma prevalence among Turkish school children (ISAAC phases I and III). *Allergy* 2006;61(12):1448-53.
20. Ripabelli G, Tamburro M, Sammarco ML, de Laurentiis G, Bianco A. Asthma prevalence and risk factors among children and adolescents living around an industrial area: a cross-sectional study. *BMC Public Health* 2013;13:1038.
21. Lai CK, Beasley R, Crane J, Foliaki S, Shah J, Weiland S. International Study of Asthma and Allergies in Childhood Phase Three Study Group. Global variation in the prevalence and severity of asthma symptoms: phase three of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax* 2009;64(6):476-83.
22. Anderson HR, Ruggles R, Pandey KD, Kapetanakis V, Brunekreef B, Lai CK, et al. isaac phase one study group. ambient particulate pollution and the world-wide prevalence of asthma, rhinoconjunctivitis and eczema in children: phase one of the international study of asthma and allergies in childhood (ISAAC). *Occup Environ Med* 2010;67(5):293-300.
23. Pearce N, Weiland S, Keil U, Langridge P, Anderson HR, Strachan D, et al. Self-reported prevalence of asthma symptoms in children in Australia, England, Germany and New Zealand: an international comparison using the ISAAC protocol. *Eur Respir J* 1993;6(10):1455-61.
24. Venn A, Lewis S, Cooper M, Hill J, Britton J. Questionnaire study of effect of sex and age on the prevalence of wheeze and asthma in adolescence. *BMJ* 1998;316(7149):1945-46.
25. Bröms K, Norbäck D, Eriksson M, Sundelin C, Svärdsudd K. Effect of degree of urbanisation on age and sex-specific asthma prevalence in Swedish preschool children. *BMC Public Health* 2009;9:303.
26. Odhiambo JA, Ng'ang'a LW, Mungai MW, Gicheha CM, Nyamwaya JK, Karimi F, et al. Urban-rural differences in questionnaire-derived markers of asthma in Kenyan school children. *Eur Respir J* 1998;12(5):1105-12.
27. Shpakou A, Brożek G, Stryzhak A, Neviartovich T, Zejda J. Allergic diseases and respiratory symptoms in urban and rural children in Grodno Region (Belarus). *Pediatr Allergy Immunol* 2012;23(4):339-46.
28. Lin RS, Sung FC, Huang SL, Gou YL, Ko YC, Gou HW, et al. Role of urbanization and air pollution in adolescent asthma: a mass screening in Taiwan. *J Formos Med Assoc* 2001;100(10):649-55.
29. Kim YK, Baek D, Koh YI, Cho SH, Choi IS, Min KU, et al. Outdoor air pollutants derived from industrial processes may be causally related to the development of asthma in children. *Ann Allergy Asthma Immunol* 2001;86(4):456-60.
30. Thirty-one source has been reshaped as "Lai C. Asthma epidemiology in Asia-Pacific. Christchurch (NZ): Asthma Asia Forum. Auckland, New Zealand, 2002".
31. Keeley DJ, Neill P, Gallivan S. Comparison of the prevalence of reversible airways obstruction in rural and urban Zimbabwean children. *Thorax* 1991;46(8):549-53.